







50°C
→

Section to be amplified







A SNP



DNA
sample 1



DNA
sample 2






β -Galactosidase

The enzyme that splits lactose into glucose and galactose. Coded by a gene ([lacZ](#)) in the [lac operon](#) of Escherichia coli.




PUC is a family of plasmids that have an ampicillin resistance gene and more importantly a *lacZ* gene. A functional *lacZ* gene will produce the protein β - galactosidase. Bacterial colonies in which β - galactosidase is produced, will form blue colonies in the presence of the substrate 5 - bromo - 4 - chloro - 3 - indolyl - β - D - galactoside or as it is more commonly referred to, X-gal.




Analytical Biochemistry **285**, 1–15 (2000)

FIG. 1. Enzymatic function of β -galactosidase in cleaving indicator substrates. β -gal cleaves β -D-galactoside containing substrates with a diverse range of aglycone groups, targeting between the glycosyl oxygen and anomeric carbon as indicated (scissors). Substrates shown indicate commonly used indicators for assays on β -gal function on plates (X-Gal) or for liquid assay by measure of fluorescence (MU-Gal or MUG) or color (ONPG). Top left, X-Gal is 5-bromo-4-chloro-3-indolyl- β -D-galactoside, and when cleaved and oxidized produces the insoluble dye 5-bromo-4-chloro-3-hydroxyindole-2-pyridone, as described previously (22). Right panel, top, yeast colonies expressing β -gal and exposed to X-Gal (right half) or the closely related compound Magenta-Gal (left half, see Biosynth, Inc., or Diagnostic Chemicals Limited). Middle left, MUG is methylumbelliferyl- β -D-galactoside, and when cleaved by β -gal produces the fluorescent product methylumbelliferone (first described in (102)). Right panel, middle, shows yeast lysates expressing β -gal exposed to MUG, under long-wave UV. Bottom left, PNPG and ONPG are closely related nitrophenol- β -D-galactosides with similar assay properties, e.g., (103), whose cleavage releases the yellow product nitrophenol (right panel, bottom); PNPG is shown.

Green Fluorescent Protein (GFP)





The **green fluorescent protein (GFP)** is a protein from the jellyfish *Aequorea victoria* that fluoresces green when exposed to blue light.




GFP Rats




Luciferase



Luciferase is a generic name for enzymes commonly used in nature for bioluminescence. The name itself is derived from *Lucifer*, which means *light-bearer*. The most famous one is firefly luciferase from the firefly *Photinus pyralis*. In luminescent reactions, light is produced by the oxidation of a luciferin (a pigment), sometimes involving Adenosine triphosphate (ATP). The rates of this reaction between luciferin and oxygen are extremely slow until they are catalyzed by luciferase, often mediated by the presence of calcium ions (an analog of muscle contraction). The reaction takes place in two steps:



Luciferase




Dim Light



Bright Light




Promega Luciferase Assay





Lipid


- ***Lipids*** are naturally occurring molecules from plants or animals that are soluble in nonpolar organic solvents.
- Lipid molecules contain large hydrocarbon portion and not many polar functional group, which accounts for their solubility behavior.





A saturated fatty acid
(palmitic acid)





A *cis* unsaturated fatty acid
(linolenic acid)




Stearic acid, an 18-carbon saturated fatty acid









A sphingomyelin



Membrane lipid



Lipid bilayer




Liposome


Properties of cell membranes:


- Cell membranes are composed of a fluid like phospholipid bilayer.
- The bilayer incorporates cholesterol, proteins, and glycolipids.
- Small nonpolar molecules cross by diffusion through the lipid bilayer.
- Small ions and polar molecules diffuse through the aqueous media in protein pores.
- Glucose and certain other substances cross with the aid of proteins without energy input.
- Na^+ , K^+ , and other substances that maintain concentration gradients inside and outside the cell cross with expenditure of energy and the aid of proteins.


- Small ions and polar molecules diffuse through the aqueous media in protein pores.
- Glucose and certain other substances cross with the aid of proteins without energy input.
- Na^+ , K^+ , and other substances that maintain concentration gradients inside and outside the cell cross with expenditure of energy and the aid of proteins.






EXTRACELLULAR FLUID











Energy and Biochemical Reactions


- Reactions in living organisms are similar to reactions in a chemical laboratory.
- Spontaneous reactions, those are favorable in the forward direction, release free energy and the energy released is available to do work.
- Spontaneous reactions , also known as *exergonic* reactions, are the source of our biochemical energy.
- Products of exergonic reactions are more stable than the reactants and the free energy change ΔG has a negative value.

- *Photosynthesis* in plants, converts CO_2 and H_2O to glucose plus O_2 which is the reverse of oxidation of glucose. The sun provides the necessary external energy for photosynthesis (686 kcal of free energy per mole of glucose formed).



- The *mitochondria* is often called the cell's power plants. Within the mitochondria, small molecules are broken down to provide the energy for an organism and also the principle energy carrying molecule adenosine triphosphate (ATP) is produced.





ATP and Energy Transfer

- Adenosine triphosphate (ATP) transport energy in living organisms.
- ATP has three $-\text{PO}_3^-$ groups.
- Removal of one of the $-\text{PO}_3^-$ groups from ATP by hydrolysis produces adenosine diphosphate (ADP). Since this reaction is an exergonic process, it releases energy.
- The reverse of ATP hydrolysis reaction is known as phosphorylation reaction. Phosphorylation reactions are endergonic.

- Biochemical energy production, transport, and use all depends on the ATP \leftrightarrow ADP interconversions.






TABLE 21.1 Free Energies of Hydrolysis of Some Phosphates





Compound Name	Function	ΔG (kcal/mol)
Phosphoenol pyruvate	Final intermediate in conversion of glucose to pyruvate (glycolysis)—stage 2, Figure 21.5	−14.8
1, 3-Bisphosphoglycerate	Another intermediate in glycolysis	−11.8
Creatine phosphate	Energy storage in muscle cells	−10.3
ATP (→ ADP)	Principal energy carrier	−7.3
Glucose 1-phosphate	First intermediate in breakdown of carbohydrates stored as starch or glycogen	−5.0
Glucose 6-phosphate	First intermediate in glycolysis	−3.3
Fructose 6-phosphate	Second intermediate in glycolysis	−3.3

- A few enzymes continuously cycle between their oxidized and reduced






Coenzyme	As Oxidizing Agent	As Reducing Agent
Nicotinamide adenine dinucleotide	NAD^+	NADH/H^+
Nicotinamide adenine dinucleotide phosphate	NADP^+	NADPH/H^+
Flavin adenine dinucleotide	FAD	FADH_2
Flavin mononucleotide	FMN	FMNH_2



The citric acid cycle

The Electron-Transport Chain and ATP Production

- Electron transport chain: The series of biochemical reactions that passes electrons from reduced coenzymes to oxygen and is coupled to ATP formation. The electrons combine with the oxygen we breathe and with hydrogen ions from their surrounding to produce water.
- Electron transport involves four enzyme complexes held in fixed positions within the inner membrane of mitochondria and two electron carriers move from one complex to another.





- Pathway of electrons in electron transport

- **ATP Synthesis**


- ADP is converted to ATP by a reaction between ADP and hydrogen phosphate ion. This is both an oxidation and phosphorylation reaction. Energy released in the electron transport chain drives this reaction forward.

Vedio: <http://www.iubmb-nicholson.org/swf/ATPSynthase.swf>







Central Dogma




Recombinant DNA







Restriction Enzyme



AluI and **HaeIII** produce blunt ends

BamHI **HindIII** and **EcoRI** produce "sticky" ends











Life

- Replication: reproduction
- Function: catalytic functions
- RNA world:
- Virus is not alive

Virus



Virus Reproduction



- Eukaryotic cells are about 1000 times larger than bacteria cells and also have a membrane enclosed nucleus containing their DNA, and several other internal structures known as organelles.





Fig 21.3 A generalized eukaryotic cell.



•Schematic showing the cytoplasm, with its components (or *organelles*), of a typical animal cell. Organelles: (1) nucleolus (2) nucleus (3) ribosome (4) vesicle (5) rough endoplasmic reticulum (6) Golgi apparatus (7) cytoskeleton (8) smooth endoplasmic reticulum (9) mitochondria (10) vacuole (11) cytosol (12) lysosome (13) centriole.


nucleus




nucleolus




nuclear envelope




ER




Golgi




lysosomes




plasma membrane




cytoplasm




centrosomes




mitochondria



microtubules




actin




with friendly permission of Jeremy Simpson and Rainer Pepperkok

A Busy Factory




A cell can be thought of as a "factory," with different departments each performing specialized tasks.

The Plasma Membrane




Cell Membrane




Characteristic diffusivities		
Particle	Typical size	Diffusion constant
Solute ion	10^{-1} nm	$2 \times 10^3 \mu\text{m}^2/\text{s}$
Small protein	5 nm	$40 \mu\text{m}^2/\text{s}$
Virus	100 nm	$2 \mu\text{m}^2/\text{s}$
Bacterium	$1 \mu\text{m}$	$0.2 \mu\text{m}^2/\text{s}$
Mammalian/human cell	$10 \mu\text{m}$	$0.02 \mu\text{m}^2/\text{s}$

The Nucleus



The cell factory contains a large inventory of blueprints dating all the way to its founding. Some of these blueprints are out of date, and some are for parts and products that are no longer made. Part of your job would entail sorting through everything, finding the correct blueprints, copying them, and sending the copies out to the assembly line at the correct time.

Nucleus




•In cell biology, the **nucleus** is a membrane-enclosed organelle found in most eukaryotic cells. It contains most of the cell's genetic material, organized as multiple long linear DNA molecules in complex with a large variety of proteins such as [histones](#) to form chromosomes. The genes within these chromosomes make up the cell's nuclear genome. The function of the nucleus is to maintain the integrity of these genes and to control the activities of the cell by regulating gene expression.

In cell biology, the **nucleolus** (plural *nucleoli*) is a "sub-organelle" of the cell nucleus, which itself is an organelle. A main function of the nucleolus is the production and assembly of ribosome components


Nuclear pores

Nuclear pores, which provide aqueous channels through the envelope, are composed of multiple proteins, collectively referred to as nucleoporins. The pores are 100 nm in total diameter; however, the gap through which molecules freely diffuse is only about 9 nm wide, due to the presence of regulatory systems within the center of the pore. This size allows the free passage of small water-soluble molecules while preventing larger molecules, such as nucleic acids and proteins, from inappropriately entering or exiting the nucleus. These large molecules must be actively transported into the nucleus instead. The nucleus of a typical mammalian cell will have about 3000 to 4000 pores throughout its envelope




Nuclear localizing sequence (NLS)

- A **nuclear localizing sequence (NLS)** is an amino acid sequence which acts like a 'tag' on the exposed surface of a protein. This sequence is used to confine the protein to the cell nucleus through the **Nuclear Pore Complex** and to direct a newly synthesized protein into the nucleus via its recognition by cytosolic nuclear transport receptors. Typically, this signal consists of a few short sequences of positively charged lysines or arginines. Typically the NLS will have a sequence (NH₂)-Pro-Pro-Lys-Lys-Lys-Arg-Lys-Val-(COOH).



The Ribosomes and the ER




Ribosomes, the workers that build proteins, are manufactured by the nucleolus. They consist of two separate subunits: a large, lower subunit and a small, upper subunit. Ribosomes attach to the rough ER. Now let's take a look at how final processing occurs

The cell has its own assembly line and workers. Within the cytoplasm is a series of large, flattened membranes that fold back and forth on each other and have a very large surface area. This collection of membranes is called the **ENDOPLASMIC RETICULUM**, or **ER**.


Ribosome

A **ribosome** is a small, dense organelle in cells that assembles proteins. Ribosomes are about 20nm in diameter and are composed of 65% ribosomal RNA and 35% ribosomal proteins (known as a [Ribonucleoprotein](#) or RNP). It translates messenger RNA (mRNA) to build a polypeptide chain (e.g., a protein) using amino acids delivered by Transfer RNA (tRNA). It can be thought of as a giant enzyme that builds a protein from a set of genetic instructions. Ribosomes can float freely in the cytoplasm (the internal fluid of the cell) or bound to the endoplasmic reticulum, or to the nuclear envelope.



Endoplasmic Reticulum

The **endoplasmic reticulum** or **ER** is an organelle found in all eukaryotic cells that is an interconnected network of tubules, vesicles and [cisternae](#) that is responsible for several specialized functions: Protein translation, folding, and transport of proteins to be used in the cell membrane (e.g., [transmembrane receptors](#) and other integral membrane proteins), or to be secreted ([exocytosed](#)) from the cell (e.g., digestive [enzymes](#)); sequestration of calcium; and production and storage of [glycogen](#), [steroids](#), and other [macromolecules](#).^[1] The endoplasmic reticulum is part of the endomembrane system. The basic structure and composition of the ER membrane is similar to the plasma membrane.




Rough endoplasmic reticulum

- The surface of the rough endoplasmic reticulum is studded with protein-manufacturing [ribosomes](#) giving it a "rough" appearance. But it should be noted that these ribosomes are not resident of the endoplasmic reticulum incessantly. The ribosomes only bind to the ER once it begins to synthesize a protein destined for sorting. The membrane of the rough endoplasmic reticulum is continuous with the outer layer of the nuclear envelope. Although there is no continuous membrane between the rough ER and the Golgi apparatus, membrane bound vesicles shuttle proteins between these two compartments. The rough endoplasmic reticulum works in concert with the Golgi complex to target new proteins to their proper destinations







Smooth endoplasmic reticulum


- The smooth endoplasmic reticulum has functions in several metabolic processes, including synthesis of lipids, metabolism of carbohydrates and calcium concentration, and attachment of receptors on cell membrane proteins. It is connected to the nuclear envelope. Smooth endoplasmic reticulum is found in a variety of cell types (both animal and plant) and it serves different functions in each. It consists of tubules and vesicles that branch forming a network. In some cells there are dilated areas like the sacs of rough endoplasmic reticulum. The network of smooth endoplasmic reticulum allows increased surface area for the action or storage of key enzymes and the products of these enzymes. The smooth endoplasmic reticulum is known for its storage of calcium ions in muscle cells.

The Golgi Apparatus



This animation illustrates how secretion vesicles, arising from the Golgi, fuse with the PM and dump the contents of their lumen outside of the cell. Note that the membrane of the SV turns insideout and becomes new PM.

-  Lipid Bilayer (SV)
-  Lipid Bilayer (PM)
-  Transmembrane Protein
-  Peripheral Protein (inside)
-  Peripheral Protein (outside)
-  Soluble Proteins





The Golgi apparatus is analogous to the finishing and packing room in a factory. Once the ribosome finishes manufacturing a protein in the rough ER, the protein needs to be prepared for use or export. Special enzymes will trim off any extra amino acids, and then the unfinished protein moves through channels in the smooth ER.

Golgi apparatus

The **Golgi apparatus** (also called the **Golgi body**, **Golgi complex**, or **dictyosome**) is an organelle found in typical eukaryotic cells. It was identified in 1898 by the Italian physician Camillo Golgi and was named after him. The primary function of the Golgi apparatus is to process and package macromolecules synthesised by the cell, primarily proteins and lipids. The Golgi apparatus forms a part of the endomembrane system present in eukaryotic cells.




Mitochondria



Like our factory's power plant, mitochondria and chloroplasts transform one form of energy to another. Remember that nearly all the energy used by living things on Earth comes from the Sun. This section discusses how energy is made available for cell processes.

Mitochondrion

- In cell biology, a **mitochondrion** is a membrane-enclosed organelle, found in most eukaryotic cells. Mitochondria are sometimes described as "cellular power plants," because they convert NADH and NADPH into energy in the form of ATP via the process of oxidative phosphorylation. A typical eukaryotic cell contains about 2,000 mitochondria, which occupy roughly one fifth of its total volume. Mitochondria contain DNA that is independent of the DNA located in the cell nucleus. According to the endosymbiotic theory, mitochondria are descended from free-living prokaryotes.







Figure 1


The main roles of the nucleolus are to synthesize rRNA and assemble ribosomes

The main function of the cell nucleus is to control gene expression and mediate the replication of DNA during the cell cycle

Lysosomes


- **Lysosomes** are organelles that contain digestive enzymes (acid [hydrolases](#)). They digest excess or worn out organelles, food particles, and engulfed viruses or bacteria. The membrane surrounding a lysosome prevents the digestive enzymes inside from destroying the cell. Lysosomes fuse with vacuoles and dispense their enzymes into the vacuoles, digesting their contents. They are built in the Golgi apparatus. The name *lysosome* derives from the [Greek](#) words *lysis*, which means dissolution or destruction, and *soma*, which means body. They are frequently nicknamed "suicide-bags" or "suicide-sacs" by cell biologists due to their role in autolysis.

Lysosomes



Lysosomes are responsible for the breakdown and absorption of materials taken in by the cell. Often, a cell engulfs a foreign substance through **ENDOCYTOSIS**, another form of active transport. During endocytosis, the cell membrane puckers up, forms a pouch around materials outside the cell, and pinches off to become a vesicle. If the contents need to be destroyed, lysosomes combine with the vesicle and release their enzymes.

Lysosome




Vedio <http://highered.mcgraw-hill.com/olc/dl/120067/bio01.swf>


Vesicle

In cell biology, a **vesicle** is a relatively small and enclosed compartment, separated from the [cytosol](#) by at least one lipid bilayer. If there is only one lipid bilayer, they are called *unilamellar* vesicles; otherwise they are called *multilamellar*. Vesicles store, transport, or digest cellular products and waste.

This biomembrane enclosing the vesicle is similar to that of the plasma membrane. Because it is separated from the cytosol, the intravesicular environment can be made to be different from the cytosolic environment. Vesicles are a basic tool of the cell for organizing metabolism, transport, enzyme storage, as well as being chemical reaction chambers. Many vesicles are made in the Golgi apparatus, but also in the endoplasmic reticulum, or are made from parts of the plasma membrane.





Cytoskeleton



The eukaryotic cytoskeleton.
Actin filaments are shown in red,
microtubules in green, and the
nuclei are in blue.



Actin



- **Actin** is a globular structural, 42 kDa, [protein](#) that polymerizes in a helical fashion to form **actin filaments** (or **microfilaments**). These form the cytoskeleton, a three-dimensional network inside the eukaryotic cell. Actin filaments provide mechanical support for the cell, determine its shape, and enable movement of the cell through [lamellipodia](#), [filopodia](#), or [pseudopodia](#). Actin filaments, along with myosin, have an essential role in muscular contraction. In the [cytosol](#), actin is predominantly bound to ATP, but can also bind to ADP. An ATP-actin complex polymerizes faster and dissociates slower than an ADP-actin complex.


Lamellipodia

- The **lamellipodium** is a cytoskeletal actin projection on the mobile edge of the cell. It contains a two-dimensional actin mesh; the whole structure pulls the cell across a substrate. Within the lamellipodia are ribs of actin called microspikes, which, when they spread beyond the lamellipodium frontier, are called filopodia (Small, et al, 2002). The lamellipodium is born of actin nucleation in the plasma membrane of the cell (Alberts, et al, 2002) and is the primary area of actin incorporation or microfilament formation of the cell. Lamellipodia range from 1 μ m to 5 μ m in breadth and are approximately 0.2 μ m thick. Lamellipodia are found primarily in very mobile cells, crawling at a speeds of 10-20 μ m/minute over epithelial surfaces..
- The tip of the lamellipodium is the site where exocytosis occurs in migrating mammalian cells as part of their clathrin-mediated endocytic cycle.




Filopodia

The **filopodia** are slender cytoplasmic projections, similar to [lamellipodia](#), which extend from the leading edge of migrating cells. They contain actin filaments cross-linked into bundles by actin-binding proteins, e.g. fimbrin. Filopodia form focal adhesions with the substratum, linking it to the cell surface. A cell migrates along a surface by extending filopodia at the leading edge. The filopodia attach to the substratum further down the migratory pathway, then contraction of stress fibres retracts the rear of the cell to move the cell forwards.



Focal adhesion



- In cell biology, '**Focal Adhesions**' are specific types of large macromolecular assemblies through which both mechanical force and regulatory signals are transmitted. More precisely, **FAs** can be considered as sub-cellular macromolecules that mediate the regulatory effects (e.g. cell anchorage) of extracellular matrix (ECM) adhesion on cell behavior.



Extra Cellular Matrix

The ECM's main components are various [glycoproteins](#), [proteoglycans](#) and [hyaluronic acid](#). In most animals, the most abundant glycoproteins in the ECM are collagens.

ECM also contains many other components: proteins such as fibrin, [elastin](#), [fibronectins](#), [laminins](#), and [nidogens](#), and minerals such as [hydroxylapatite](#), or fluids such as blood plasma or serum with secreted free flowing [antigens](#).




Integrin


An **integrin**, or **integrin receptor**, is an integral membrane protein in the plasma membrane of cells. It plays a role in the attachment of a cell to the extracellular matrix (ECM) and to other cells, and in signal transduction from the ECM to the cell. There are many types of integrin, and many cells have multiple types on their surface. Integrins are of vital importance to all metazoans, from humans to sponges.

Schematic drawing of a typical integrin dimer

Arrow shows the region where an I domain is inserted in some α subunits. Not all α subunits are posttranslationally cleaved. Internal disulphide bonds within subunits are not shown. Dark blue regions in the head segment of the α subunit represent homologous repeats. Those with the EF-hand consensus sequence are marked with red circles to denote binding sites for divalent metal ion.



Endocytosis



Endocytosis

- Phagocytosis is the process by which cells ingest large objects, such as cells which have undergone apoptosis, bacteria, or viruses. The membrane folds around the object, and the object is sealed off into a large [vacuole](#) known as a [phagosome](#).
- Pinocytosis is a synonym for endocytosis. This process is concerned with the uptake of solutes and single molecules such as proteins.
- Receptor-mediated endocytosis is a more specific active event where the cytoplasm membrane folds inward to form coated pits. These inward budding vesicles bud to form cytoplasmic vesicles.

Endocytosis pathways

- Macropinocytosis is the invagination of the cell membrane to form a pocket which then pinches off into the cell to form a vesicle filled with extracellular fluid (and molecules within it). The filling of the pocket occurs in a non-specific manner. The vesicle then travels into the [cytosol](#) and fuses with other vesicles such as [endosomes](#) and [lysosomes](#).
- Clathrin-mediated endocytosis is the specific uptake of large extracellular molecules such as proteins, membrane localized receptors and ion-channels. These receptors are associated with the cytosolic protein clathrin which initiates the formation of a vesicle by forming a crystalline coat on the inner surface of the cell's membrane.
- [Caveolae](#) consist of the protein caveolin-1 with a bilayer enriched in cholesterol and glycosphingolipids. Caveolae are flask shaped pits in the membrane that resemble the shape of a cave (hence the name caveolae). Uptake of extracellular molecules are also believed to be specifically mediated via receptors in caveolae.

Gene Therapy

- Gene therapy is a technique for correcting defective genes responsible for disease development. Researchers may use one of several approaches for correcting faulty genes:
 - **A normal gene may be inserted into a nonspecific location within the genome to replace a nonfunctional gene. This approach is most common.**
 - **An abnormal gene could be swapped for a normal gene through homologous recombination.**
 - **The abnormal gene could be repaired through selective reverse mutation, which returns the gene to its normal function.**
 - **The regulation (the degree to which a gene is turned on or off) of a particular gene could be altered.**


How Gene Therapy Works?

- In most gene therapy studies, a "normal" gene is inserted into the genome to replace an "abnormal," disease-causing gene. A carrier molecule called a vector must be used to deliver the therapeutic gene to the patient's target cells. Currently, the most common vector is a virus that has been genetically altered to carry normal human DNA. Viruses have evolved a way of encapsulating and delivering their genes to human cells in a pathogenic manner. Scientists have tried to take advantage of this capability and manipulate the virus genome to remove disease-causing genes and insert therapeutic genes.
- Target cells such as the patient's liver or lung cells are infected with the viral vector. The vector then unloads its genetic material containing the therapeutic human gene into the target cell. The generation of a functional protein product from the therapeutic gene restores the target cell to a normal state.

Gene Delivery

- Transfection- the delivery of foreign molecules such as DNA and RNA into eukaryotic cells
- Naked DNA is not suitable for in-vivo transport of genetic materials-> degradation by serum nucleases
- Ideal gene delivery system
 - Biocompatible
 - Non-immunogenic
 - Stable in blood stream
 - Protect DNA during transport
 - Small enough to extravagate
 - Cell and tissue specific

Endocytosis



Endocytosis

- Phagocytosis is the process by which cells ingest large objects, such as cells which have undergone apoptosis, bacteria, or viruses. The membrane folds around the object, and the object is sealed off into a large [vacuole](#) known as a [phagosome](#).
- Pinocytosis is a synonym for endocytosis. This process is concerned with the uptake of solutes and single molecules such as proteins.
- Receptor-mediated endocytosis is a more specific active event where the cytoplasm membrane folds inward to form coated pits. These inward budding vesicles bud to form cytoplasmic vesicles.

Endocytosis pathways

- Macropinocytosis is the invagination of the cell membrane to form a pocket which then pinches off into the cell to form a vesicle filled with extracellular fluid (and molecules within it). The filling of the pocket occurs in a non-specific manner. The vesicle then travels into the [cytosol](#) and fuses with other vesicles such as [endosomes](#) and [lysosomes](#).
- Clathrin-mediated endocytosis is the specific uptake of large extracellular molecules such as proteins, membrane localized receptors and ion-channels. These receptors are associated with the cytosolic protein clathrin which initiates the formation of a vesicle by forming a crystalline coat on the inner surface of the cell's membrane.
- [Caveolae](#) consist of the protein caveolin-1 with a bilayer enriched in cholesterol and glycosphingolipids. Caveolae are flask shaped pits in the membrane that resemble the shape of a cave (hence the name caveolae). Uptake of extracellular molecules are also believed to be specifically mediated via receptors in caveolae.

Endocytic pathway in mammalian cells




Figure 1 Multiple portals of entry into the mammalian cell. The endocytic pathways differ with regard to the size of the endocytic vesicle, the nature of the cargo (ligands, receptors and lipids) and the mechanism of vesicle formation.




Figure 2 Cargo-stimulated signalling pathways induce uptake by phagocytosis and caveolae. **a**, Fc receptors on the surface of macrophages are activated by immunoglobulin- γ molecules bound to a bacterium. A signalling cascade that involves Rac, Cdc42 and downstream kinases triggers actin rearrangements, protrusion of the membrane around the bacterium, and its engulfment into a phagosome. **b**, Albumin binds to and presumably clusters its receptor, gp60, in caveolae to activate $G_{i\alpha}\beta/\gamma$ and Src kinases, triggering caveolae endocytosis.






Figure 3 Core components of the machinery driving clathrin-mediated endocytosis. Clathrin triskelions, composed of three clathrin heavy chains (CHC) and three tightly associated light chains (CLC), assemble into a polygonal lattice, which helps to deform the overlying plasma membrane into a coated pit. Heterotetrameric AP2 complexes are targeted to the plasma membrane by the α -adaptin subunits, where they mediate clathrin assembly through the $\beta 2$ -subunit, and interact directly with sorting motifs on cargo molecules through their $\mu 2$ subunits. Dynamin is a multidomain GTPase that is recruited to the necks of coated pits, where it can assemble into a spiral or 'collar' to mediate or monitor membrane fission and the release of CCVs (see text for details). A subsequent uncoating reaction recycles the coat constituents for reuse.





Formation of Clathrin-Coated Vesicles




- 2500 every minute
- CCV uncoat within seconds

Clathrin-mediated endocytosis



A



B

Lipid Rafts



Caveolae



Caveolin



Phospholipid



Sphingolipid



Cholesterol

Barrier to non-viral gene delivery




Figure 1 Barriers to non-viral gene delivery

Representation of the route travelled by a non-viral gene-delivery vector carrying therapeutic DNA to the nucleus. A non-viral vector, formed by interaction of the DNA with a carrier compound, must cross the plasma membrane to enter the cell. This can be via several routes, including endocytosis-based entry (1a), direct physical entry routes, such as electroporation or ballistic delivery (1b), or direct entry via protein transduction (1c). Depending on the mode of cellular entry, the vector may become encapsulated in an endosome (2), from which it must escape (3a) or it will become degraded when the endosome fuses with a lysosome (3b). The DNA will at some point be subjected to degradation by cytosolic nucleases (4), as it traverses through the cytoplasm to reach the nucleus. Finally, the vector must undergo nuclear transport (5) through NPCs embedded in the NE in order to gain access to the nucleoplasm. Once in the nucleus, the DNA may (6a) or may not (6b) need to be uncoated, depending upon the vector used, before it can ultimately be transcribed (7).

NLS-mediated nuclear import




Figure 2 NLS-mediated nuclear import pathways

In classical nuclear import, the NLS found in cargo bound for the nucleus is recognized by the Imp α subunit of the Imp α/β heterodimer (1). However, there are also many examples where Imp β or one of its many homologues can mediate nuclear import or cargo proteins independently of Imp α (2). In both cases, transient interactions between the Imp β and the nucleoporin proteins that line the NE-embedded NPCs mediate translocation into the nucleus. Once inside, RanGTP binds to Imp β (3), releasing Imp α and the cargo into the nucleoplasm (4a and 4b). RanGTP itself is then recycled back to the cytoplasm (5), where it is converted into its RanGDP state (not shown). An animated version of this Figure can be found at <http://www.BiochemJ.org/bj/406/0185/bj4060185add.htm>

Transfection

