

DM physics at the ILC

Shigeki Matsumoto (Kavli IPMU)

***Physics perspectives at the ILC experiment
on the systematical search for the WIMP DM.***

Current status of the ILC project in Japan

May 27, 2013

Asking Science Council of Japan (SCJ) to deliberate the ILC project from an academic point of view.

Dec 24, 2013

\$0.5M for investigatory study was approved. The same amount was approved for the following year.

May 01, 2014

An academic expert committee (ACE) was established under MEXT.

By Mar 31, 2016 (Extendable)
MEXT will receive a report from ACE.

LCC (LC collaboration) activities including ILC TDR completion, etc.

Sep 30, 2013 (SCI)

A report was submitted to MEXT.

- Discussing scientific significance.
- Verifying TDR in details.
- Investigating economical ripple effect, technology transfer, etc.

Academic expert committee

May 08, 2014: 1st Meeting of ACE

Nov 05, 2014: 2nd Meeting of ACE

Apr 10, 2015: 3rd Meeting of ACE
[Interim report came out]

Jun XX, 2015: 4th Meeting of ACE

...

Physics perspectives at the ILC

Why the electroweak symmetry is broken at $O(100)\text{GeV}$?

I. Precise measurements of the Higgs boson and the top quark

New physics@ $O(1)\text{TeV}$ predicts the deviation of Higgs couplings (and top couplings) from SM predictions at $O(1)\%$ level. Precise measurement of m_t is also important for BSMs behind the EWSB.

II. Direct and indirect searches for new particles (new phenomena)

BSM behind often predicts new particles at around $O(1)\text{TeV}$. Among those, the most promising candidate is WIMP dark matter!

Model-dependent way

Merit: Well-defined framework / Connection to a big picture.

Demerit: Many possibilities of BSMs / No signals so far at the LHC.

Model-independent way

Merit: Investigating the entire concept of the WIMP dark matter.

Demerit: Framework tends to be complicated.

Searching for WIMP dark matter

It is convenient to categorize WIMP DM candidates according to their quantum numbers: **Spin and Weak isospin**. (EM and color charges = 0)

Weak isospin of the WIMP DM:

1. Singlet-like case:

WIMP DM is an (almost) singlet under SM gauge interactions $SU(3)_c \times SU(2)_L \times U(1)_Y$. Typical examples are the bino, the singlino, etc.

2. Multiplet-like case:

WIMP DM is an (almost) neutral component of a $SU(2)_L$ multiplet. Typical examples are the Higgsino, the wino, the minimal DM, etc.

3. Mixed case:

WIMP DM is a mixture of different $SU(2)_L$ representations due to the EWSB. Typical examples are Bino-Higgsino dark matter, etc.

We consider these 3 cases with the dark matter being a fermion.

DM cosmology puts an upper limit on the WIMP DM mass in any case!

$$\Omega_{\text{DM}} h^2 = \Omega_{\text{TH}} h^2 + \Omega_{\text{NT}} h^2 = 0.11 \text{ with } \Omega_{\text{TH}} h^2 \propto (m_{\text{DM}})^2 \text{ and } \Omega_{\text{NT}} h^2 > 0$$

WIMP DM search (Mixed case)

Characteristics:


- DM has a sizable coupling to the Higgs boson (DM-DM-Higgs).
- Several non-colored particles exist in the same mass scale of DM.
- The mass of the DM will be in between the EW and TeV scales.

The WIMP DM will be discovered at DM direct detection experiments or the LHC at first. (This case is gradually being excluded by experiments so far, but there are still parameter regions evading these limits.)




The role of the ILC is to study its property in details.

Production of EM charged (non-colored) particles decaying to DM



Signal: 4-jets (from WW) + Emiss

Energy dist of reconstructed Ws



[Asano, S.M. et al. 1106.1932]


Property is determined in detail!

WIMP DM search (Multiplet-like case)


Characteristics:

- DM is highly degenerate with EM charged $SU(2)_L$ partners.
- Physics of the DM is governed mainly by $SU(2)_L$ gauge interaction.
- The mass of the DM can be as high as $O(1)TeV$ when $\Omega_{NT} h^2$ is small.

Production of $SU(2)_L$ partners



Signal: $\gamma + E_{miss} + \text{soft tracks}$




$m_{DM} < 0.5 s^{1/2} - 20GeV$ is explored.
Property is determined in detail.
[Berggren et al. 1307.3566]

Corrections to SM processes



Signal: Interference with SM one



Sys1. e:0.2%, μ :0.15%, b:0.5%, c:1.0%
Sys2. e:0.4%, μ :0.30%, b:1.0%, c:2.0%
 $m_{DM} < s^{1/2} / 2 + 200GeV$ explored.

WIMP DM search (Multiplet-like case)



TeV scale thermal WIMP (DM relic abundance only from $\Omega_{\text{TH}} h^2$):

- When DM is from a $SU(2)_L$ doublet, its mass is predicted to be 1TeV.
- When DM is from a $SU(2)_L$ triplet, its mass is predicted to be 3TeV.



None of current & near future collider experiments can access the DM. Only DM indirect detection experiments have a possibility to detect it.

A hint of the TeV scale WIMP from the AMS-02 anti-p/p data:



If this is true, CLIC or 100TeV (with a perfect detector) is needed.

WIMP DM search (Singlet-like case)

Characteristics:

- No renormalizable interaction with SM particles (DM-DM-SM).
- The mass of the DM is within a very natural region, say EW scale.



We have to consider an EFT involving higher dimensional operators.

$$\mathcal{L}_{F_0} = \mathcal{L}_{\text{SM}} + \frac{1}{2} \bar{\chi} (i\not{\partial} - M_\chi) \chi + \sum_{n,i} \frac{c_i^{(n)}}{\Lambda^{n-4}} \mathcal{O}_i^{(n)}$$

$$\begin{aligned} \mathcal{O}_S^{(5)} &= (\bar{\chi}\chi)|H|^2, \\ \mathcal{O}_H^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(H^\dagger i\overleftrightarrow{D}_\mu H), \\ \mathcal{O}_Q^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{Q}\gamma_\mu Q), \\ \mathcal{O}_U^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{U}\gamma_\mu U) \\ \mathcal{O}_D^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{D}\gamma_\mu D) \\ \mathcal{O}_L^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{L}\gamma_\mu L) \\ \mathcal{O}_E^{(6)} &= (\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{E}\gamma_\mu E) \end{aligned}$$

With CP invariance and flavor blindness in the DM sector, we have 7 independent SM gauge invariant operators.




We have to consider the EFT with all 'c_i' being treated as independent couplings.

[S. M., Mukhopadhyay, Tsai, 1405.1859]



Clarifying parameter regions consistent with current limits and go to ILC issues.

WIMP DM search (Singlet-like case)



Limits considered:

Relics abundance: $\Omega_{\text{DM}} h^2 < 0.1$

Direct detection: SI, SD-p, SD-n

Collider (LEP): Mono- γ , Γ_Z

Collider (LHC): Mono-j, Γ_h




Shape of the region (twin peaks) is determined mainly by $\Omega_{\text{DM}} h^2$.


Hunting grounds of the ILC

$$\begin{aligned} &(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{L}\gamma_\mu L) \\ &(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{E}\gamma_\mu E) \end{aligned}$$

$$(\bar{\chi}\gamma^\mu\gamma_5\chi)(H^\dagger i\overleftrightarrow{D}_\mu H)$$



Signal: mono- γ search ($\gamma + E_{\text{miss}}$)



Summary

The ILC will offer a unique way to **search for the WIMP dark matter**.

Mixed case:

WIMP DM will be discovered by DM detect detection experiments or the LHC. The role of the ILC is to investigate its property in detail.

Multiplet-like case:

When WIMP DM is not heavy, it will be discovered by the production of EM charged $SU(2)_L$ partners or the deviation from SM prediction at $e^-e^+ \rightarrow f\bar{f}$ processes. Property of DM can also be investigated.

Singlet-like case:

Wide parameter regions will be covered by the ILC via the mono- γ search. This case is now regarded as the most important one, for it predicts the WIMP DM at EW scale and none of other experiments can touch it. The analysis based on our method is now being a Key project with ILC experimentalists, and the result will be reflected to the design of ILC detectors.

Backup slides

Precise Higgs measurements

Please refer also to K. Fujii's talk slides in PHENO 2014

I = 1.15ab^{-1} @ 250GeV \oplus 1.6ab^{-1} @ $500(550)\text{GeV}$ & II = $\oplus 2.5\text{ab}^{-1}$ @ 1TeV

	Δm_h	Γ_h	κ_t	κ_b	κ_τ	κ_c	κ_Z	κ_W	κ_g	κ_γ	λ
I.	30MeV	2.5%	7.8(2.3)%	0.8%	1.2%	1.5%	0.5%	0.6%	1.2%	4.5(1.7)%	46%
II.	---	2.3%	1.9(1.7)%	0.7%	0.9%	1.0%	0.5%	0.6%	0.9%	2.4(0.8)%	13%

↳ [$s^{1/2} = 550\text{GeV}$ I/O 500GeV] [LHC-ILC synergy] ↓


ILC can determine couplings in a model independent way

- $\sigma \times \text{BR}$ measured in each mode
- σ measured from recoil mass, namely, $e^-e^+ \rightarrow H Z \rightarrow X \mu^- \mu^+$.
- Total width of H obtained by $e^-e^+ \rightarrow \nu\nu H \rightarrow \nu\nu W^-W^+$. ($s^{1/2} > 350\text{GeV}$ is crucial.)



Coupling determinations!

An implication to SUSY scenario



$m_h = 126\text{GeV}$ & several constraints are already included in the analysis.
 $m_A = \text{a few TeV can be covered.}$


Precise top measurements

Please refer also to K. Fujii's talk slides in PHENO 2014
10fb⁻¹ each (11bins) @ 340-350GeV ⊕ 500fb⁻¹ @ 500GeV

$$\Gamma_{\mu}^{ttX}(k^2, q, \bar{q}) = ie \left\{ \gamma_{\mu} \left(\tilde{F}_{1V}^X(k^2) + \gamma_5 \tilde{F}_{1A}^X(k^2) \right) + \frac{(q - \bar{q})_{\mu}}{2m_t} \left(\tilde{F}_{2V}^X(k^2) + \gamma_5 \tilde{F}_{2A}^X(k^2) \right) \right\}$$


Δm_t	$\Delta \Gamma_t$	\tilde{F}_{1V}^{γ}	\tilde{F}_{1V}^Z	\tilde{F}_{1A}^Z	\tilde{F}_{2V}^{γ}	\tilde{F}_{2V}^Z
100MeV	32MeV	± 0.002	± 0.002	± 0.006	± 0.001	± 0.002

Stability of the Higgs potential



Boundary condition $\lambda(m_{pl}) = 0$?
Precise measurements of m_t !

An implication to CHMs, etc.



Precise top measurements can be used to discriminate NPMs!

WIMP DM search (Singlet-like case) [Next step]

No renormalizable interactions between the WIMP DM and SM particles.


→ Considering higher dimensional interactions up to dimension six.

[7 SM gauge invariant Ops with CP/Flavor inv. e.g. $(c/\Lambda^2)(\bar{\chi}\gamma_\mu\gamma_5\chi)(\bar{e}_R\gamma^\mu e_R)$]

This EFT may not work for physics of energetic colliders such as ILC.

→ Considering two different UV theories reproducing the EFT.

UV⁺ ⊃ Heavy Z₂-even particles




$\Lambda \rightarrow \infty$

EFT ⊃ Higher dimensional Ops.




$\Lambda \rightarrow \infty$

UV⁻ ⊃ Heavy Z₂-odd particles



[S.M., Mukhopadhyay, Tsai, 1047.1859, etc.]



EFT, UV⁺, UV⁻ are used for collider signals and we accept the most conservative one.